# **Curly Hair Tarantula**

Brachypelma albopilosum OR Tliltocatl albopilosus





## **Physical Characteristics**

- <u>Physical description:</u> Long, curly hairs called setae with barbs all over its body. Hair is densest
  near their hind legs. Eight segmented legs and eight eyes. Body separated into a cephalothorax
  and an abdomen. Hair can be 3-4 inches long. Two pedipalps on the front of their
  cephalothorax.
- <u>Coloration:</u> Bronze body with light brown to black hairs. The hair can also have a grey or pink color. Juveniles are paler than the adults.
- <u>Sexual dimorphism:</u> Males are lighter in color.

# Size

- Length: 5-5.5 in (12.7-14 cm)
- Weight: Females can weigh up to 2 ounces.

# Lifespan

- Wild: Males 4 years, Females 8-10 years
- Zoos: Males 6 years, Females 30 years

## **Behaviors**

- Activity Cycle: Nocturnal
- <u>Lifestyle:</u> Terrestrial, sleep in burrows during the day. They do not travel far when they come out of their burrows to feed.
- Vocalizations: Hissing to scare predators
- Social Structure: Solitary
- Will form webs that line their burrows to protect their eggs.

# Diet

- Wild: Small vertebrates like lizards, frogs, and mice, as well as large insects like crickets, roaches, and grasshoppers.
- Zoo: Crickets

# Reproduction

- Incubation Period: 8-12 weeks
- Parenting: Mothers take care of juveniles for a few weeks.
- Clutch Size: 300-500 golf ball-sized eggs.
- Breeding Behaviors: Males will find a female's burrow and tap their leg on the ground. If the female agrees to mate, the male creates a sperm web and gives it to the female. The male may then be allowed to leave or may be eaten by the female.

#### **Habitat and Range**

- <u>Habitat:</u> Found in burrows in deciduous forest, evergreen rainforests, and grasslands
- Range: Nicaragua, Costa Rica, and Honduras

## Conservation

- <u>IUCN Status:</u> Not Listed on IUCN, but CITES required permits to trade this species.
- Contributing factors: Habitat loss, pet trade, and people killing them out of fear.

#### Classification

• Kingdom: Animalia



<u>Phylum</u>: Arthropoda<u>Class:</u> Euchelicerata

• Order: Araneae

• <u>Family</u>: Theraphosidae

• Genus: Tliltocatl or Brachypelma

• Species: T. albopilosus or B. albopilosum

# **Interesting Facts**

- Predators include multiple species of birds, snakes, lizards, wasps, and other spiders.
  - The tarantula hawk wasp is a parasitoid that uses curly haired tarantulas as hosts. A parasitoid is an animal that is a parasite only in their larval stage.
- Another name for this tarantula is the Honduran Curly Hair.
- Recent research has indicated that there are two different species of curly haired tarantulas. Tliltocatl albopilosus are from Nicaragua and Brachypelma albopilosum come from Honduras.
- Their hair does not just protect them, it also helps them sense movement in their environment.